arXiv:0808.3280v2 [cond-mat.mtrl-sci] 2 Nov 2008

Photon blockade and nonlinear effects for a quantum dot strongly coupled to a semiconductor microcavity

F. Bello* and D. M. Whittaker
Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, United Kingdom
(Dated: November 2, 2008)

Our model comprehensively simulates modern nanoscale semiconductor microcavities incorporating cavity quantum electrodynamics within both the weak and strong coupling regimes, using on-resonant laser excitation and nonresonant excitation due to a wetting layer. For weak coupling, the most significant effect is photon antibunching with nonresonant emission. We investigate how the antibunching characteristics change as the cavity finesse is increased towards the strong coupling regime. Antibunching can also be observed in a strongly coupled system with resonant excitation, using the photon blockade mechanism which has been demonstrated in atom systems. We calculate what cavity parameters are required to observe this effect. Experimentally these studies are equivalent to nonlinear pump probe measurements, where a strong pump, either resonant or nonresonant, is used to excite the coupled system, and the resulting state is characterized using a weak, resonant probe beam.

PACS numbers:

Currently, a single atom trapped within a cavity is the standard model for cavity quantum electrodynamics(CQED) with the semiconductor analogue of this being a quantum dot(QD) trapped within a dielectric cavity. In this paper, we investigate theoretically CQED effects that may be observed using a coupled QD/cavity system while taking into consideration the current standards of cavity finesse and coupling strength(g_c). In particular, we choose to focus on antibunching effects which occur as a direct consequence of the interaction between cavity field and QD.[7] The Jaynes-Cummings dressed state model forms the eigenstates of such a system in which the energy difference between any two eigenstates is unique due to their dependence on the number of particles in the system.[8] This subsequently permits the system to absorb one resonant photon at a time, thus creating a photon blockade and therefore a single photon emitter. This type of device is useful in many genres of physics, for instance, quantum cryptography or quantum computation; fields that both require the entanglement and emission of single particles within discrete energy levels.

Quantitatively antibunching is demonstrated by computing a second order correlation function at zero time delay($g^{(2)}(\tau = 0)$) that is less than one, a purely quantum mechanical effect, with the ideal value for perfectly antibunched light being zero. Previous work has been conducted on isolated quantum dots[9,10] and more recently QDs or atoms within cavities[4] testing their ability to produce antibunched light. The most successful photon blockades showed a $g^{(2)}(0)$ approaching values of 0.1 for isolated electrically pulsed dots as well as measurements reported to be as low 0.13±.11 within strongly coupled atom/cavity systems.[1] This research makes direct comparisons with each case in which we present data for the strong and weak coupling regimes and concentrate on various schemes that can be used to pump a QD trapped within a microcavity, namely, via a wetting layer or resonant excitation with a laser. With on-resonant pumping one has the benefit of being able to directly create an exciton in the QD hence reducing emission from off-resonant energy levels which could occur through incoherent pumping as the electron relaxes into the dot. This also avoids the delayed capturing of the electron or hole and the consequential ionization of the QD. In addition, cavities have the known advantages of aiding in the collection efficiency of photons as well as promoting recombination dynamics.

The separation between weak and strong coupling regimes is found by making a comparison of the ratio between cavity and dot decay rates ($\gamma_{cav}, \gamma_{dot}$) and the size of the energy splitting[4] for which one wants

$$\sqrt{N_p} g_c \gg \gamma_{cav}, \gamma_{dot} \tag{1}$$

with the number of particles, N_p, in the system increasing the energy separation between successive pairs of dressed state levels. At the moment decay rates for a typical microcavity are usually two orders of magnitude larger than that of a QD which are presented in the formalism for completeness although considered to be negligible in the final calculation. Strong coupling is not easily obtained in current QD/microcavity systems. Atoms tend to offer a bigger ratio for strong coupling vs. decoherence time being greater than 8:1[1] compared to less than 1:1 for quantum dots.[3] Dots however have the benefit of not only being stable components of the solid state structure, but restricting particle movement in all three dimensions, while in principle, the manufacturability of quantum dot structures will allow them to emit at any optical frequency requested. In order to better achieve strong coupling within a QD/microcavity structure one requires a small modal volume to decrease the decay rate and increase the cavity's Q factor, $\frac{\lambda}{\Delta\lambda}$. Micropillar cavities draw particular interest in the field due to their manufacturable control over the modal volume as well as the strong overlap attainable between QD and cavity mode. Most recently strong coupling effects such as the anticrossing of dot and cavity energies have been observed within micropillars having a modal volume of 0.43μm and a Q factor between 10,000-20,000.[2,4]

For the purpose of calculating antibunching effects we want to explicitly write the correlation function in terms of the definitions for the first($i = 1$) and second($i = 2$) order Green's functions given as $G^{(i)}(0) = tr(\rho(t)b_1^\dagger b_2^\dagger ... b_i^\dagger b_i ... b_2 b_1)$. Here we represent the cavity field using second quantization operators, $b^\dagger$ and (b), creating or annihilating a cavity photon, respectively, with $\rho(t)$ being the steady state solution to the density matrix. The normalized second order correlation function is then defined as

$$g^{(2)}(0) = \frac{G^{(2)}(0)}{|G^{(1)}(0)|^2} = \frac{\sum_n n(n-1)P_n}{\bar{n}^2} \tag{2}$$

where P_n is the probability of finding the system with n photons. The summation over the photon number, n, comes from using a Foch or number state representation as a basis with the denominator equal to the average number of photons in the system defined as $\bar{n} = \sum nP_n$. Physically this calculation equivocates to a simultaneous readout taken by two detectors within the Hanbury Brown-Twiss experiment and computing the average over the fluctuations in output intensity(I), yielding an alternative yet equivalent definition,

$$g^{(2)}(0) - 1 = \frac{\langle (I - \langle I \rangle)^2 \rangle}{\langle I \rangle^2}. \tag{3}$$

The ideal scenario, corresponding to having $g^{(2)}(0) = 0$, occurs when the detectors cannot simultaneously count a photon, hence insuring that one has a consistent single photon emitter.

Special attention is placed on the type of pumping, either coherent or incoherent, that one would use in order to perform pump-probe type experiments. Firstly, we look at incoherent excitation of the dot via a wetting layer since it is the mechanism more commonly used. When one does so, there are certain conditions which perpetuate stronger antibunching for weak coupling. For example, background interference is diminished if the pumping of the QD is reduced as well as the subsequent loss of the dot's spontaneous emission rate into a cavity mode due to the Purcell effect, which vanishes as g_c goes to zero or the quality factor diminishes. A 3-level system is used to represent the dot and wetting layer as depicted in figure 1. The ground state is the source for excitations within the system, hence the wetting layer is coherently excited from the ground state of the dot with field strength Γ. Pumping of the dot is subsequently modeled via the incoherent decay rate, γ_w, from the wetting layer to the excited state of the dot where an electron can pair with a hole in the valence band. The hamiltonian given in the rotating frame is

$$\begin{aligned} H &= \hbar\delta_d\sigma_2 + \hbar\delta_w\sigma_3 + \hbar\delta_c b_j^\dagger b_j \\ &+ \hbar g_c(\sigma_{12}^+ b_j + b_j^\dagger \sigma_{12}^-) + \Gamma(\sigma_{13}^+ + \sigma_{13}^-), \end{aligned} \tag{4}$$



Figure 1: Schematic representing a 3-level system consisting of a wetting layer(w) and quantum dot(ω_d) which is coupled to a cavity mode(ω_c) via the coupling strengh(g_c). The wetting layer is coherently driven by an on-resonant laser(Γ) whilst the dot is being incoherently pumped directly from the wetting layer to the excited state of the dot.



Figure 2: $g^{(2)}(0)$ shown as a function of coupling over decay for incoherent pumping of the dot via a wetting layer with γ_{cav} kept constant. Pumping from the wetting layer increases in order from lowest to highest going from a dotted to a more progressively solid line. The coupling g_c is varied for different pumping rates from the wetting layer to the excited state of the dot. A crossover appears around the weak to strong coupling transition at which point the affect of pump strength on antibunched light is inverted.

where we have used a product basis consisting of an expansion of the pauli spin matrices into three dimensions in order to describe the dot and wetting layer with its ground state energy set equal to zero. Again, the Foch states are used to describe the mode making use of the ladder operators b and $b^\dagger$. As defined analogous to reference[6], $\sigma_{rs}^+ = \sigma_{rs}^{-*} = \frac{1}{2}(S_x^{rs} + iS_y^{rs})$ are fermionic operators that describe interactions between levels r and s, and act on either the dot's ground(1) and excited(2) states along with the wetting layer's(3). $\sigma_i = |i\rangle\langle i|$ is a level shift operator whose expectation value reveals the population of energy level 'i' while $\delta_{d,c,w}$ are the detunings for the dot(d), cavity(c), and wetting layer (w) respectively, relative to the laser frequency taken to be an on-resonant cw π-pulse. We look to solve for the density matrix, given below, of this coupled system in order to determine its dynamics.

$$\begin{aligned}\dot{\rho} &= -\frac{i}{\hbar}[H_s,\rho] - \frac{\gamma_{cav}}{2}[b^\dagger b\rho(t) + \rho(t)b^\dagger b - 2b\rho(t)b^\dagger] \\ &- \frac{\gamma_{dot}}{2}[\sigma_{12}^+\sigma_{12}^-\rho(t) + \rho(t)\sigma_{12}^+\sigma_{12}^- - 2\sigma_{12}^-\rho(t)\sigma_{12}^+] \\ &- \frac{\gamma_w}{2}[\sigma_{23}^+\sigma_{23}^-\rho(t) + \rho(t)\sigma_{23}^+\sigma_{23}^- - 2\sigma_{23}^-\rho(t)\sigma_{23}^+] \qquad (5)\end{aligned}$$

Explicitly shown are decay terms ($\gamma_{cav(dot,w)}$) taking into consideration the interaction of the microcavity with a surrounding reservoir. These terms are well known and derived using Markovian statistics. Antibunching effects are calculated solving for the second order Green's function presented in equation 2. As shown in figure 2, a crossing occurs around the weak to strong coupling transition that is dependent on the cavity's decay rate which we keep constant in the present study and only vary g_c. It may appear that the curves cross at the same point but under close inspection they *expectedly* do not since the strong coupling condition varies following the ratio given in equation 1. In the strong coupling regime, the best antibunching happens for the highest pumping from the wetting layer(solid curve) in direct contrast to weak coupling results. The crossover could be understood by utilizing the form of $g^{(2)}(0)$ as it appears in the number state basis given in equation 2. We therefore could write the difference of the second order correlation function between one'higher'(H) and 'lower'(L) pumping cases appearing in figure 2 as,

$$g_D^{(2)}(0) = \sum_{n=2}^{\infty} n(n-1)\frac{P_n^H}{\bar{n}_H^2}(1 - \frac{P_n^L}{P_n^H}\frac{\bar{n}_H^2}{\bar{n}_L^2}). \qquad (6)$$

Using the incoherent pumping scheme proposed, the distribution of light is a complicated function depending on the intricate balance between the ratios of γ_w, γ_{cav}, g_c, and Γ, however the general lineshape does resemble that predicted for antibunched light.[13] From the numerics we see that for all parameters studied the following relationship holds for $n \geq 2$,

$$(\bar{n}_H^2 - \bar{n}_L^2) > 0 > P_n^L - P_n^H. \qquad (7)$$

In addition, within the strong coupling regime we have the relation that

$$|\bar{n}_H - \bar{n}_L| > |P_n^L - P_n^H|, \qquad (8)$$

thus guaranteeing that the last term in parenthesis in equation 6 is greater than one, therefore yielding a negative value for $g_D^{(2)}(0)$. As figure 2 shows for this regime, this result states that a higher pumping rate leads to a higher average number of photons in the cavity which will therefore yield a lower $g^{(2)}(0)$. Equation 8 suggests that under strong coupling the average number of photons dominates the second order correlation function and hence by examining equation 2, the denominator will dictate photon antibunching where increasing(decreasing) $\bar{n}$ would decrease(increase) bunching.



Figure 3: $g^{(2)}(0)$ shown as a function of detuning(det) over coupling(g_c) for various coupling parameters increasing sequentially from the present day standards for QD/cavity structures with $g_c/\gamma_c = 1$(black) to that for atom/cavity structures with $g_c/\gamma_c = 8$(blue). Here, the driving field has directly excited the cavity mode. Antibunching effects ($g^{(2)}(0) < 1$) occur when the driving field is off-resonance by roughly the energy level splitting $\hbar g_c$ at which point $g^{(2)}(0)$ tends toward zero creating a single photon emitter. Highly off-resonance we return to the expected value of one for a coherent driving field . The weakly coupled case(black) remains circa the classical barrier of one for a coherent laser field.

As we move into the weak coupling regime the relationship in Equation 8 is no longer valid. There is a lower probability of the dot emitting a photon into the cavity due to the smaller coupling to decay ratio as well as the loss of the Purcell effect. The population is still significant enough to be able to emit consistently with $\bar{n}$ reaching as high as 0.25, however in this regime $\bar{n}_H \approx \bar{n}_L$ allowing the ratio P_n^L/P_n^H in equation 6 to dominate the last term of $g_D^{(2)}(0)$ forcing it to be positive. At very low coupling the correlations, $G^2(0)$, and the avergage number of photons, $\bar{n}$, both tend toward zero yet are comparable to each other, hence we still calculate a $g^{(2)}(0)$ although emission from the cavity is deficient. When $g_c = 0$, no photons are entering the cavity($\bar{n} = 0$) at which point calculation of the second order correlation is ineffectual. The coupling parameter alone regulates the transition rate from exciton to cavity photon. By looking at any curve in figure 2, all for which γ_w is kept constant, when the coupling is increased more photons are allowed to enter the cavity as they are fed into the dot via the wetting layer, therefore increasing the bunching in the system. The curve begins to level off as the rate at which the wetting layer populates the dot is overcome by the transition rate from QD to cavity. At this point the strength of the coupling has little affect on the number of photons in the cavity if the dot isn't being sufficiently pumped at a comparable rate. In the weak coupling regime, the model is essentially analogous to an isolated quantum dot with effective antibunching but



Figure 4: $g^{(2)}(0)$ shown as a function of detuning(det) over coupling(g_c) for various coupling parameters increasing sequentially from the present day standards for QD/cavity structures with $g_c/\gamma_c = 1$(black) to that for atom/cavity structures with $g_c/\gamma_c = 8$(blue). The driving field in this case is meant to directly excite the dot which in turn populates the cavity. Prominent antibunching affects occur in the region when detuning equals coupling strength. Off-resonance one does not return to a coherent driving field but towards a squeezed state.



Figure 5: Comparison of $g^{(2)}(0)$ for coherent excitation of the dot vs. cavity while varying g_c. Presented is the case when detuning is equal to the energy level splitting, $\hbar g_c$. As coupling increases the lines tend to meet as $g^{(2)}(0)$ goes towards zero for perfect antibunching .

without the benefits of being enclosed in the microcavity. A value comparable to the previously mentioned results for isolated QDs of approximately 0.1 is shown but in actuality if one reduces the incoherent pumping rate the correlation function will tend towards zero.

We now choose to look at the pumping scheme for on-resonant excitation of a cavity mode with a laser. For the case of a coherently pumped cavity we have the following Hamiltonian given in the rotating frame as,

$$\begin{aligned} H &= \hbar\delta_d\sigma_z + \hbar\delta_c b_j^\dagger b_j \\ &+ \hbar g_c(\sigma^+ b_j + b_j^\dagger\sigma^-) + \Gamma(b^\dagger + b). \end{aligned} \tag{9}$$

The Pauli spin matrices are used to describe the 2-level dot with the subscripts 1,2 on $\sigma^{+(-)}$ dropped. Second quantization is used once again for the cavity mode which is excited by a classical driving field. By changing the operators in the Γ term from $b^\dagger(b)$ to $\sigma^+(\sigma^-)$, we will be able to consider the laser field coherently exciting the QD as well. The master equation is given by the following expression,

$$\begin{aligned} \dot\rho &= -\frac{i}{\hbar}[H_s,\rho] - \frac{\gamma_{cav}}{2}[b^\dagger b\rho(t) + \rho(t)b^\dagger b - 2b\rho(t)b^\dagger] \\ &- \frac{\gamma_{dot}}{2}[\sigma^+\sigma^-\rho(t) + \rho(t)\sigma^+\sigma^- - 2\sigma^-\rho(t)\sigma^+] \end{aligned} \tag{10}$$

and as expected, applying a conventional Jaynes-Cummings model using the classical driving field, we attain analogous results for that of an atom coupled to a cavity mode.[1] Figure 3 shows that when the light field is on resonant with the energy splitting(g_c) we should witness the strongest antibunching of photons creating a photon blockade effect. The blue curve represents the present day atom/cavity standard for a coupling to decay ratio of about 8:1. Within the present day standards for the QD/cavity system (1:1 ratio), the system shows no antibunching effects(black curve). As detuning increases we return to the expected value of one for a coherent driving field. Going through the transition of weak to strong coupling(black to red curve), one sees a general increase in antibunching creating a more reliable single photon emitter with the best results for the strongest coupling. At zero detuning a strong bunching of photons will occur on the detectors due to correlations between the pump and cavity/dot mode.[14] Additional peaks in figure 3 as well as figure 4 are due to the occupation of higher energy levels in the dressed state model, namely a two-particle eigenstate with an energy splitting of $\pm g_c/\sqrt{2}$.

Figure 4 considers direct excitation of the dot via use of a coherent driving field. Presently, distributed Bragg reflectors (DBR mirrors) are commonly used to restrict a photon's movement along the vertical axis of a microcavity pillar structure. Confinement in the lateral direction due to strong internal reflection is dependent on the angle of incidence to the pillar's external wall as well as the index of refraction of the medium. Relative to the angle of incidence, ongoing research is looking into whether the exciton could be directly created by the laser in lieu of excitation via an absorbed cavity mode. Antibunching effects are shown to be more easily obtainable within this method with the value of the correlation function going well below 0.1. Off-resonance from the coupling strength, the $g^{(2)}(0)$ function no longer returns to that of a coherent laser field, but rather it is a squeezed state still emitting single photons. This improvement in antibunching could be explained due to the use of a QD to populate the cavity instead of a coherent field. The two-level dot acts as a natural antibunching device for light permitting only one photon to enter the cavity at a time. With a sufficient coupling to cavity decay ratio, the photon will leave the cavity before a saturated dot reemits another into it allowing particles to bunch in the system.

A comparison of $g^{(2)}(0)$ under both pumping schemes is presented in figure 5. Direct excitation of the QD consistently shows a $g^{(2)}(0) < 1$ with perfect antibunching seen at a significantly lower coupling to decay rato than that for direct exciation of a cavity mode. Results such as these are highly noteworthy in regards to today's capabilities for manufacturing and manipulating strongly coupled cavity/QD systems as well as the present understanding of what characteristics are needed if such a system will be able to produce antibunched light. It has been demonstrated that within the weak coupling regime, populating your system via a wetting layer could produce a lower $g^{(2)}(0)$ than in the strongly coupled regime comparable to present day standards, however the advantages of cavity containment and on-resonant excitation are lost. In addition, in order to perform a successful photon blockade, we have shown that coherent driving fields demand strongly coupled systems when exciting a cavity mode, however the possibility of directly exciting the quantum dot laxes this requirement and produces a more strongly antibunched squeezed light field.

Acknowledgments

We would like to thank Almut Biege and Daniele Sanvitto for useful conversations and insight on the subject presented.

---

[*] Electronic address: f.bello@shef.ac.uk

[1] K. M. Birnbaum, A. Boca, R. Miller, A. D. Boozer, T. E. Northup, and H. J. Kimble, Nature **436**, doi:10.1038 (2005).

[2] A. Löffler, J. P. Reithmaier, G. Sek, C. Hofmann, S. Reitzenstein, M. Kamp, and A. Forchel, Appl. Phys. Lett. **86**, 111105 (2005).

[3] T. Yoshie, A. Scherer, J. Hendrickson, G. Khitrova, H. M. Gibbs, G. Rupper, C. Ell, O. B. Shchekin, and D. G. Deppe, Nature **432**, doi:10.1038 (2004).

[4] D. Press, S. Götzinger, C. Reitzenstein, A. Hofmann, M. Löffler, A. Kamp, Y. Forchel, and Y. Yamamoto, Phys. Rev. Lett. **98**, 117402 (2007).

[5] P. Meystre and M. Sargent III, *Elements of Quantum Optics, 3rd edn.* (1998).

[6] D. Suter, *The Physics of Laser-Atom Interactions* (1997).

[7] F. P. Laussy, M. M. Glazov, A. Kavokin, D. M. Whittaker, and G. Malpuech, Phys. Rev. B. **73**, 115343 (2006).

[8] E. Jaynes and F. Cummings, Proc. IEEE **51**, 89 (1963).

[9] Z. Yuan, B. E. Kardynal, R. M. Stevenson, A. J. Shields, C. J. Lobo, K. Cooper, N. S. Beattie, D. A. Ritchie, M. Pepper, Science **295**, 102 (2002).

[10] R. J. Young, D. J. P. Ellis, R. M. Stevenson, A. J. Bennett, P. Atkinson, K. Cooper, D. A. Ritchie, and A. J. Shields, Proc. IEEE **95**, 9 (2007).

[11] M. Fox, *Quantum Optics*, (2006).

[12] A. Kavokin, J. J. Baumberg, G. Malpuech, and F. P. Laussy, *Microcavities*, (2007).

[13] H. J. Carmichael, *Statistical Methods in Quantum Optics 2: non-classical fields*, (2008).

[14] P. R. Rice and H. J. Carmichael, IEEE J. Quant. Elect. **24**, 7 (1988).